

Extraction Post-op Instructions for Parents

If your child has had 1 or more teeth taken out – "wiggled":

- 1. The gauze needs to stay in place with biting pressure for 15 minutes.
- 2. Give your child the appropriate dose of children's Tylenol, Motrin or Advil when you take the gauze out. Your child should only need this for approximately 12 to 24 hours. If pain persists beyond 48 hours, call our office.
- 3. Your child should eat only soft food for one to two days nothing crunchy and nothing too hot or cold because the area may be a little sensitive. Encourage plenty of liquids (water, soups, juices, etc.). A regular diet can be started as tolerated by your child.
- NO SPITTING OR DRINKING THROUGH A STRAW OR SIPPY CUP! This can cause the bleeding to start again.
- A clean mouth heals faster. Gentle brushing around the extraction site can be started immediately.
- 6. Activity may need to be limited; sometimes a nap is a good idea.
- 7. Swelling after an extraction is not uncommon and is no cause for concern. Apply an ice pack for
 15 minutes on and 15 minutes off as needed in the 24 hours following tooth removal.
- 8. Your child's cheek, lip and tongue will be numb for approximately 2 hours. As this area "wakes up" and the numbness goes away, it may feel funny. Please be very careful that your child does not bite at his/her cheek or pick at this area. Only liquid or non-chewing foods during the 2 hours: smoothies or milkshakes (no straw!), yogurt, pudding, apple sauce are allowed. Please avoid foods with seeds for the next three days.